Cowlitz County Diabetes Statistics, 2003



 Prevalence
 Cowlitz County
 Washington State

 (adults 18+ years)
 Percent (95% CI)¹ Number of adults
 Percent (95% CI)¹ Number of adults

 5.4 (3.5, 8.3)
 4,230
 6.6 (6.2, 7.1)
 303,283

Diabetes Prevention & Control Program P.O. Box 47836 Olympia, WA 98504-7836 (360) 236-3617

Risk Factors	Cowlitz County	Washington State
(adults 18+ years without diabetes)	Percent (95%CI) ¹	Percent (95%CI) ¹
Overweight or obese ²	66.2 (59.0, 72.8)	56.8 (55.8, 57.8)
Insufficient physical activity ³	31.4 (25.3, 38.2)	35.8 (34.9, 36.8)
Smoke cigarettes	23.1 (18.1, 29.0)	19.8 (19.0, 20.6)
History of high blood pressure ⁴	25.3 (19.9, 31.5)	20.8 (20.1, 21.5)
History of high cholesterol ⁴	23.7 (18.3, 30.1)	31.1 (30.2, 32.1)

_	Number of	Age-adjusted rate per 10,000 total		Crude rate per 10,000 people with	
Hospitalizations ^⁵	hospitalizations	population (95% CI) ¹		diabetes	
(with complications of diabetes)	Cowlitz County	Cowlitz County	Washington State	Cowlitz County	Washington State
Diabetes as first listed diagnosis	87	8.9 (7.1, 11.1)	9.6 (9.4, 9.9)	202.4	191.3
Diabetes as any listed diagnosis	1,367	133.3 (126.3, 140.6)	118.4 (117.5, 119.3)	3,180.6	2,293.6
with coronary heart disease	455	43.7 (39.8, 48.0)	36.3 (35.8, 36.8)	1,058.7	693.1
→ with stroke	78	7.6 (6.0, 9.5)	10.1 (9.9, 10.4)	181.5	190.9
→ with diabetic ketoacidosis	38	4.2 (3.0, 5.8)	3.9 (3.8, 4.1)	88.4	78.9
with lower extremity amputation	28	2.7 (1.8, 4.0)	1.9 (1.8, 2.0)	65.1	160.7

Deaths ⁵	Number of deaths	Age-adjusted rate per 100,000 total population (95% CI) 1		Crude rate per 100,000 people with diabetes	
	Cowlitz County	Cowlitz County	Washington State	Cowlitz County	Washington State
Diabetes as underlying cause	29	27.3 (18.3, 39.6)	26.0 (24.7, 27.4)	674.7	494.4
Diabetes as any cause	113	107.7 (88.7, 129.9)	77.1 (74.9, 79.4)	2,629.2	1,460.9

Please see http://www.doh.wa.gov/cfh/diabetes/publications/Face%20Page.pdf for a description of data sources and methods used to develop these county profiles (includes estimates of total diabetes population for each county that are used to compute the hospitalization and death rates per people with diabetes).

¹Data presented as: percentage or rate (95% confidence interval).

²Overweight=body mass index (BMI) 25.0-29.9 kg/m², obese=BMI 30.0+ kg/m².

³Insufficient physical activity=reported level of physical activity during leisure time or at work does not meet Centers for Disease Control and Prevention (CDC) recommendations of moderate-intensity activity for 30+ minutes on 5+ days/week or vigorous-intensity activity for 20+ minutes on 3+ days/week

⁴Told by doctor, nurse, or other health professional had high blood pressure and/or high cholesterol.

⁵Rates are not computed for counts <5. Rates based on fewer than 20 deaths or hospitalizations are likely to be unstable or imprecise.